

SPECIMEN LABEL



PROFESSIONAL INSECT CONTROL® ROACH KILLER BAIT GEL

SPECIMEN LABEL

# ROACH KILLER BAIT GEL (3-Pack)

## FOR USE IN COMMERCIAL, INDUSTRIAL AND RESIDENTIAL AREAS

### ACTIVE INGREDIENT:

Fipronil\*: [5-Amino-1-(2,6-dichloro-4-(trifluoromethyl)phenyl)-4-(1,R,S)-(trifluoromethyl)sulfinyl)-1H-pyrazole-3-carbonitrile] ..... 0.01%

OTHER INGREDIENTS: ..... 99.99%

\*CAS NO. 120068-37-3

Contains: 3 plastic dispensers

## KEEP OUT OF REACH OF CHILDREN

### CAUTION: SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Use only in areas not easily accessible to children and pets.

Product may stain textiles and clothing.

### PRECAUTIONARY STATEMENTS:

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Keep gel away from foods and food contact surfaces. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling.

**FIRST AID:** IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists. IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

**PHYSICAL AND CHEMICAL HAZARDS:** Do not use this product in or on electrical equipment where a possibility of shock hazard exists.

**STORAGE AND DISPOSAL:** Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a cool, dry area out of reach of children.

**DISPOSAL:** Wrap used pre-filled plastic reservoirs in several layers of newspaper and discard in trash.

If you have questions regarding the use of this product, please call 1-800-426-6228.

Maxforce website address: [www.maxforce.com](http://www.maxforce.com)

### DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

MAXFORCE® FC Professional Insect Control® Roach Killer Bait Gel may be used as a spot or crack and crevice treatment for the indoor and outdoor control of cockroaches in residential areas and the non-food/non-feed areas of institutional, warehousing and commercial establishments, including warehouses, restaurants, food processing plants, supermarkets, hospitals, nursing homes, motels, hotels, schools, laboratories, computer facilities, sewers, "aircraft," buses, boats/ships, trains, pet shops and zoos. Non-food/non-feed areas are areas such as garbage rooms, lavatories, entries, and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling). This product may also be used in food/feed areas of food/feed handling establishments as outlined below.

Application rate is dependent on level of infestation and species to be controlled. Bait can be applied as spots or as a bead in cracks and crevices. The lower application rate should be used for low to moderate infestations or re-treatments, and the higher rate for more severe infestations or where the population occurs in inaccessible areas, such as in walls. Gel placements should be at or near harborage or aggregation areas, such as corners, areas of movement or cracks and crevices. Numerous smaller placements will provide faster control than fewer larger spots, especially for German cockroach control.

**For outdoor use:** Place bait in areas adjacent to structures where roaches may nest or breed. Apply bait to potential points of insect entry, such as eaves, sills, and expansion joints. Apply bait in protected areas whenever possible. Environmental stresses such as direct sunlight and water will reduce the residual effectiveness.

Priority treatment areas may be identified by visual inspection of such areas listed above or aided by trapping techniques or spot flushing.

Printed 12/2000

**Refillable Stations:** This product may be used in the Maxforce Refillable Bait Station or B&G Perimeter Patrol Station. Do not exceed 1 tsp. of gel per refillable station. Refillable stations should be placed every 20 to 30 feet around the perimeter of a structure. Use approximately 2 stations per side of a typical single family home.

For best results, place stations near: air conditioning units, water faucets, foundation vents and weep holes, decks, utility entryways (e.g. electrical, cable TV) and wherever target pests are frequently found. Regularly inspect all stations and replace bait as needed for continual control.

Position Refillable Bait Stations adjacent to structure in areas where they will not be damaged by lawn equipment such as mowers and trimmers. When placing the station in turf remove sod layer and place station base plate against the soil.

If the station is not equipped with a sticker to identify the contents, apply a sticker to the outside or inside of the station.

#### Applications in Food/Feed Handling Establishments: Application within food/feed areas of food/feed handling establishments is limited to crack and crevice treatment only.

Apply with syringe applicator directly into cracks and crevices. Place applicator tip 1/2 inch into cracks, crevices, holes and other small openings where roaches may be a problem. Apply small amounts of gel (up to 1/2 gram) per spot. Do not apply bait to areas where food/feed, utensils or processing surfaces may become contaminated. Do not apply bait to areas that are routinely washed such as cracks and crevices in tops of tables, food/feed preparation and prepared food holding surfaces as bait may be removed by washing. Do not apply bait to surfaces where the temperature exceeds 130°F as bait may run. Examples include, but are not limited to, portions of stoves, ovens, grills, fume hoods, heat lamps, coffee urns, steam tables, toasters, fryers, dishwashers and hot water pipes. Care should be taken to avoid depositing gel onto exposed surfaces. If gel contacts an exposed surface, remove gel and wash exposed surface.

Food/feed areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing edible waste storage, and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas are also considered a food/feed area when food is exposed and facility is in operation. Apply bait in small amounts between different elements of construction, between equipment and floors, openings leading into voids and hollow spaces in floors, walls, ceilings, equipment legs and bases, around plumbing pipes, doors and windows, behind and under refrigerators, cabinets, sinks, closets, stoves and other equipment where cockroaches hide. During follow-up visits inspect bait placements and reapply when necessary. **Application of this product in food/feed areas of food/feed handling establishments other than as a crack and crevice treatment is not permitted.**


Keep exposed gel away from open food and food contact surfaces. Do not treat food preparation surfaces. Do not apply bait gel to areas which have been recently sprayed with insecticide, and do not spray insecticide over gel as it may cause the bait to become repellent. Do not place treatments in locations where routine cleansing operations may transfer diluted bait to food or food preparation surfaces.

**Contains: 3 plastic dispensers  
NET WT 2.1 OZ (60g) each  
TOTAL NET WT 6.3 OZ (180g)**

Distributed by & © 1998  
Maxforce Insect Control Systems  
1221 Broadway  
Oakland, CA 94612



EPA Reg. No. 64248-14  
EPA Est. No. 5813-MS-01

SPECIES / LEVEL OF INFESTATION	APPLICATION RATE (PER SQ. YD.)
<b>MODERATE INFESTATIONS OR RETREATS FOR GERMAN OR BROWN-BANDED ROACHES:</b>	1-2 SPOTS ABOUT THE SIZE OF THE GEL SYRINGE CAP OR 1 1/2" - 3" BEAD
<b>SEVERE OR HEAVY INFESTATIONS FOR GERMAN OR BROWN-BANDED ROACHES:</b>	2-4 SPOTS ABOUT THE SIZE OF THE GEL SYRINGE CAP OR 3" - 6" BEAD
<b>LARGE ROACHES SUCH AS AMERICAN, SMOKEY-BROWN OR ORIENTAL:</b>	 (Actual size syringe cap) RECOMMENDED SPOT SIZE (1/4 - 1/2 g)